


RESEARCH NOTE

Open Access



# A retrospective cross-sectional survey on nosocomial bacterial infections and their antimicrobial susceptibility patterns in hospitalized patients in northwest of Iran

Hamid Sadeghi<sup>1</sup>, Saeideh Gholamzadeh Khoei<sup>2</sup>, Mehdi Bakht<sup>3</sup>, Mohammad Rostamani<sup>3</sup>, Sara Rahimi<sup>3</sup>, Mehdi Ghaemi<sup>4</sup> and Bahman Mirzaei<sup>1,5\*</sup> 

## Abstract

**Objective:** Nosocomial infections (NIs) are known as one of the remarkable problems in all countries. This study is aimed to estimate the prevalence rate of nosocomial bacterial agents with antimicrobial susceptibility pattern in hospitalized patients. This study was conducted from April 2017 to September 2018, on 4029 hospitalized patients. We set out to recognize the commonest bacterial infections and antimicrobial susceptibility patterns of nosocomial infection.

**Results:** Of the 4029 patients, 509 (12.6%) of them were culture positive. Of these *Escherichia coli* (*E. coli*) (98.3%) and *Staphylococcus epidermidis* (*S. epidermidis*) (37.5%) were the most abundant bacterial identified in the urinary tract and bloodstream cultures respectively. Moreover, *Acinetobacter spp.* (100%) and *Pseudomonas aeruginosa* (22.2%) were the most abundant organisms detected in the respiratory system. According to susceptibility testing results, 370 (80.5%) and 264 (57.3%) in Gram-negatives and 44 (91.7%) and 35 (72.9%) in gram positives isolated strains were classified as multidrug-resistant (MDR) and extensive drug-resistant (XDR) strain respectively. On account of the high prevalence of MDR and XDR bacterial species, there is a pressing need for the expansion of new strategies on antibiotic supervision and infection control to introduce new guideline on empirical antibiotic therapy.

**Keywords:** Nosocomial infections, Bacterial isolates, Susceptibility patterns, Multi-drug resistant (MDR), Extensively-drug resistant (XDR)

## Introduction

Nosocomial infections (NIs) are known as hospital-gained infection expanding within 48–72 h after incoming [1]. A large part of morbidity and mortality in hospitals goes back to the NIs [2]. Following the increasing rate of NIs, developing in socio-economic disturbance, antimicrobial resistance, and the mortality rate are inevitable [3]. NIs happen throughout the world both

in developing and developed countries. NIs accounts for 10% in developing and 7% in developed countries [4]. Nearly 2 million people have been tangled in this matter and also would be known as a major reason for the loss of life and money [5]. Bacteria are hugely the prominent cause of NIs [6]. The most usual kinds of NIs which happen in a hospital set up are: urinary system, bloodstream infections, surgical site infections, gastroenteritis, respiratory system [7, 8]. Despite global endeavor to rein NIs during previous years, NIs still stays a widespread problem and as one of the important causes of antibiotic resistance in hospitalized patients [9]. As antibiotic resistance continues to greaten, attuned definitions with which

\*Correspondence: dr.bahman.m@gmail.com; drbahman@zums.ac.ir

<sup>1</sup> Department of Microbiology and Virology, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran

Full list of author information is available at the end of the article



© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Medical Sciences. In the current study, all ethical guidelines including Ethics and Consent to participate from the parents have been collected.

# Consent to publication

Not applicable.

# Competing interests

The authors announce that they have no difference in interest.

# Author details

<sup>1</sup> Department of Microbiology and Virology, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran. <sup>2</sup> Molecular Medicine Research Center, Hamadan University of Medical Sciences, Hamadan, Iran. <sup>3</sup> Department of Microbiology, Qazvin University of Medical Sciences, Qazvin, Iran. <sup>4</sup> Department of Anesthesiology, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran. <sup>5</sup> Department of Microbiology and Virology, School of Medicine, Mazandaran University of Medical Sciences, Mazandaran, Iran.

Received: 12 October 2020 Accepted: 26 February 2021

Published online: 09 March 2021

# References

- Vincent J-L. Nosocomial infections in adult intensive-care units. *Lancet*. 2003;361(9374):2068–77.
- Pourakbari B, Rezaizadeh G, Mahmoudi S, Mamishi S. Epidemiology of nosocomial infections in pediatric patients in an Iranian referral hospital. *J Prevent Med Hyg*. 2012;53(4).
- Organization WH: Report on the burden of endemic health care-associated infection worldwide. 2011.
- Khan HA, Baig FK, Mehboob R. Nosocomial infections: epidemiology, prevention, control and surveillance. *Asian Pac J Trop Biomed*. 2017;7(5):478–82.
- Mohajeri P, Azizkhani S, Farahani A, Norozi B. Genotyping of coa and aroA Genes of methicillin-resistant staphylococcus aureus strains isolated from nasal samples in western Iran. *Jundishapur J Microbiol*. 2016. <https://doi.org/10.5812/jjm.26460>.
- System NNIS. National Nosocomial Infections Surveillance (NNIS) System report, data summary from January 1990–May 1999, issued June 1999. *Am J Infect Control*. 1999;27(6):520–32.
- Endalafer N, Gebre-Selassie S, Kotiso B. Nosocomial bacterial infections in a tertiary hospital in Ethiopia. *J Infect Prevent*. 2011;12(1):38–43.
- Raka L, Zoutman D, Mulliqi G, Krasniqi S, Dedushaj I, Raka N, Ahmeti S, Shala M, Vishaj A, Elezi Y. Prevalence of nosocomial infections in high-risk units in the university clinical center of Kosova. *Infect Control Hosp Epidemiol*. 2006;27(4):421–3.
- Farr BM, Salgado CD, Karchmer TB, Sherertz RJ. Can antibiotic-resistant nosocomial infections be controlled? *Lancet Infect Dis*. 2001;1(1):38–45.
- Anderson DJ, Engemann JJ, Harrell LJ, Carmeli Y, Reller LB, Kaye KS. Predictors of mortality in patients with bloodstream infection due to ceftazidime-resistant *Klebsiella pneumoniae*. *Antimicrob Agents Chemother*. 2006;50(5):1715–20.
- Cosgrove SE, Sakoulas G, Perencevich EN, Schwaber MJ, Karchmer AW, Carmeli Y. Comparison of mortality associated with methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* bacteremia: a meta-analysis. *Clin Infect Dis*. 2003;36(1):53–9.
- Roberts RR, Hota B, Ahmad I, Scott RD, Foster SD, Abbasi F, Schabowski S, Kampe LM, Ciavarella GG, Supino M. Hospital and societal costs of antimicrobial-resistant infections in a Chicago teaching hospital: implications for antibiotic stewardship. *Clin Infect Dis*. 2009;49(8):1175–84.
- Ibrahim EH, Sherman G, Ward S, Fraser VJ, Kollef MH. The influence of inadequate antimicrobial treatment of bloodstream infections on patient outcomes in the ICU setting. *Chest*. 2000;118(1):146–55.
- Cohen AL, Calfee D, Fridkin SK, Huang SS, Jernigan JA, Lautenbach E, Oriola S, Ramsey KM, Salgado CD, Weinstein RA. Recommendations for metrics for multidrug-resistant organisms in healthcare settings: SHEA/HIC-PAC position paper. *Infect Control Hosp Epidemiol*. 2008;29(10):901–13.
- Hidron AI, Edwards JR, Patel J, Horan TC, Sievert DM, Pollock DA, Fridkin SK. Antimicrobial-resistant pathogens associated with healthcare-associated infections: annual summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2006–2007. *Infect Control Hosp Epidemiol*. 2008;29(11):996–1011.
- Falagas ME, Karageorgopoulos DE. Pandrug resistance (PDR), extensive drug resistance (XDR), and multidrug resistance (MDR) among Gram-negative bacilli: need for international harmonization in terminology. *Clin Infect Dis*. 2008;46(7):1121–2.
- Park YK, Peck KR, Cheong HS, Chung D-R, Song J-H, Ko KS. Extreme drug resistance in *Acinetobacter baumannii* infections in intensive care units, South Korea. *Emerg Infect Dis*. 2009;15(8):1325.
- Magiorakos A-P, Srinivasan A, Carey R, Carmeli Y, Falagas M, Giske C, Harbarth S, Hindler J, Kahlmeter G, Olsson-Liljequist B. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect*. 2012;18(3):268–81.
- Halaji M, Feizi A, Mirzaei A, Sedigh Ebrahim-Saraie H, Fayyazi A, Ashraf A, Havaei SA. The global prevalence of class 1 integron and associated antibiotic resistance in *Escherichia coli* from patients with urinary tract infections, a systematic review and meta-analysis. *Microb Drug Resist*. 2020;26(10):1208–18.
- Halaji M, Shahidi S, Atapour A, Ataei B, Feizi A, Havaei SA. Characterization of extended-spectrum  $\beta$ -lactamase-producing uropathogenic *Escherichia coli* among Iranian kidney transplant patients. *Infect Drug Resistance*. 2020;13:1429.
- Zalipour M, Esfahani BN, Halaji M, Azimian A, Havaei SA. Molecular characterization of vancomycin-resistant *Enterococcus faecalis* among inpatients at Iranian University Hospitals: clonal dissemination of ST6 and ST422. *Infect Drug Resist*. 2019;12:3039.
- Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. *Am J Infect Control*. 1988;16(3):128–40.
- Ducel G, Fabry J, Nicolle L. Prevention of hospital acquired infections: a practical guide. *Prevention of hospital acquired infections: a practical guide* 2002(Ed. 2).
- Cheesbrough M. District laboratory practice in tropical countries, part 2. Cambridge: Cambridge University Press; 2006.
- Wayne P. Clinical and laboratory standards institute. Performance standards for antimicrobial susceptibility testing. 2011.
- Zahedi M, Abounoori M, Maddah MM, Mirabi A, Sadeghnezhad R, Rezaei AA, Goli HR. Evaluation of bacterial nosocomial infections and antibiotic resistance pattern: a 2-year epidemiological surveillance study in a hospital population. *Int J Med Investig*. 2019;8(3):91–103.
- Davoudi AR, Najafi N, Shirazi MH, Ahangarkani F. Frequency of bacterial agents isolated from patients with nosocomial infection in teaching hospitals of Mazandaran University of Medical Sciences in 2012. *Caspian J Intern Med*. 2014;5(4):227.
- Bijari B, Abbasi A, Hemati M, Karabi K. Nosocomial infections and related factors in southern Khorasan hospitals. *Iran J Med Microbiol*. 2015;8(4):69–73.
- Rahimi-Bashar F, Karami P, Khaledi A, Dehghan A, Seifrabie MA, Yaghoobi MH. Evaluation of the prevalence of nosocomial infection in different wards of Be'sat hospital of Hamedan. *Avicenna J Clin Microbiol Infect*. 2018;5(2):31–5.
- Zahraei SM, Eshrafi B, Asl HM, Pezeshki Z. Epidemiology of four main nosocomial infections in Iran during March 2007–March 2008 based on the findings of a routine surveillance system. *Arch Iran Med*. 2012;15(12):0–0.
- Keihanian F, Saeidinia A, Abbasi K, Keihanian F. Epidemiology of antibiotic resistance of blood culture in educational hospitals in Rasht, North of Iran. *Infect Drug Resist*. 2018;11:1723.
- Rajabi M, Abdar ME, Rafiei H, Aflatoonia MR, Abdar ZE. Nosocomial infections and epidemiology of antibiotic resistance in teaching hospitals in south east of Iran. *Glob J Health Sci*. 2016;8(2):190.
- Zamani S, Nasiri MJ, Khoshnab BN, Ashrafi A, Abdollahi A. Evaluation of antimicrobial resistance pattern of nosocomial and community bacterial pathogens at a teaching hospital in Tehran, Iran. *Acta Medica Iranica* 2014;182–186.
- Soltani R, Ehsanpoor M, Khorvash F, Shokri D. Antimicrobial susceptibility pattern of extended-spectrum  $\beta$ -lactamase-producing bacteria causing nosocomial urinary tract infections in an Iranian referral teaching hospital. *J Res Pharm Pract*. 2014;3(1):6.